


Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.


In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

International Journal of Antimicrobial Agents

journal homepage: <http://www.elsevier.com/locate/ijantimicag>

Short communication

Detection of class 1 and 2 integrons, extended-spectrum β -lactamases and *qnr* alleles in enterobacterial isolates from the digestive tract of Intensive Care Unit inpatients[☆]

Inés Bado^a, Nicolás F. Cordeiro^a, Luciana Robino^a, Virginia García-Fulgueiras^a, Verónica Seija^b, Cristina Bazet^b, Gabriel Gutkind^c, Juan A. Ayala^d, Rafael Vignoli^{a,*}

^a Departamento de Bacteriología y Virología, Instituto de Higiene, Facultad de Medicina, Universidad de la República, Av. Alfredo Navarro 3051, CP 11600, Montevideo, Uruguay

^b Laboratorio Clínico del Hospital de Clínicas 'Dr Manuel Quintela', Facultad de Medicina, Universidad de la República, Av. Italia s/n, CP 11600, Montevideo, Uruguay

^c Cátedra de Microbiología, Facultad de Farmacia y Bioquímica, Junín 956, CP 1113, Buenos Aires, Argentina

^d Centro de Biología Molecular 'Severo Ochoa', CSIC-UAM, Consejo Superior de Investigaciones Científicas, Campus de Cantoblanco, 28049 Madrid, Spain

ARTICLE INFO

Article history:

Received 22 March 2010

Accepted 21 June 2010

Keywords:

Integrons
Transferable
Resistance
*bla*_{CTX-M}

ABSTRACT

In this study, we searched for extended-spectrum β -lactamases (ESBLs), class 1 and 2 integrons, and *qnrA*, *qnrB* and *qnrS* genes in 56 oxyimino-cephalosporin and/or ciprofloxacin-resistant enterobacterial isolates obtained from the gastrointestinal tract of patients admitted in an Intensive Care Unit in Uruguay. ESBLs were detected in 11 isolates (6 CTX-M-2, 3 CTX-M-9, 1 CTX-M-15 and 1 PER-2). *qnr* genes and integrons were detected in 5 and 24 isolates, respectively. Eight different antibiotic resistance gene cassettes were found within six different genetic arrangements. Two types of complex class 1 integrons carrying insertion sequence *ISCR1* were found, one showing *bla*_{CTX-M-2}-*orf3* and the other *qnrA1*-*ampR*. Ten of the thirteen isolates carrying class 2 integrons presented the element *IS5* inserted between *intI2* and *dfrA1*, whereas another class 2 integron lacked the internal stop codon usually present in *intI2*. This is the first report of the occurrence of *qnrA*, *qnrB* and *bla*_{CTX-M-9} in Uruguay. Dissemination of the different groups of CTX-M enzymes (i.e. groups 1, 2 and 9) appears to be a recent phenomenon in South America.

© 2010 Elsevier B.V. and the International Society of Chemotherapy. All rights reserved.

1. Introduction

The increase in antibiotic resistance is a growing problem worldwide, both in the hospital and the community setting [1,2], owing to the appearance of new antibiotic resistance genes and/or the optimisation of mechanisms responsible for their mobilisation. Amongst the new resistance genes that have gained notoriety in recent years, those coding for transferable quinolone resistance [3] as well as the pandemic expansion of extended-spectrum β -lactamases (ESBLs) of the CTX-M family [1] may be different faces of multiresistance evolution.

The occurrence in South America of *qnr* genes and of the enzyme AAC(6')-Ib-cr is a recent phenomenon [3]. Moreover, ESBLs from the CTX-M group appear to follow two distinct geographical distribution patterns; whilst in Argentina only CTX-M-2 group enzymes

have been reported [4], the occurrence of various CTX-M groups in other countries appears to be the rule [2]. In Uruguay, the presence of CTX-M-2 has already been reported [5,6].

In addition, new forms of gene acquisition by mobile genetic elements have also emerged, such as class 1 integrons linked to the insertion sequence *ISCR1* (complex integrons) [1] and class 2 integrons with a functional integrase capable of inserting new gene cassettes into their variable region [7].

Recently, we reported for the first time in Uruguay the presence of two *Escherichia coli* isolates carrying *bla*_{CTX-M-15} and *aac(6')-Ib-cr* along with a class 1 integron carrying *dfr17*-*aadA5* as gene cassettes [8].

The objectives of this study were to detect and characterise class 1 and 2 integrons, ESBLs and *qnr* genes in enterobacterial isolates obtained from Intensive Care Unit (ICU) inpatients in Uruguay.

2. Material and methods

2.1. Patients

In total, 104 patients requiring mechanical respiratory assistance were included in this study. The patients were admitted to the

[☆] This work was presented in part at the 48th Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC) and the 46th Annual Meeting of the Infectious Diseases Society of America (IDSA), 25–28 October 2008, Washington DC (abstract C2-755).

* Corresponding author. Tel.: +598 2 487 57 95; fax: +598 2 487 57 95.

E-mail address: rvignoli@higiene.edu.uy (R. Vignoli).

ICU of the University Hospital of Montevideo, Uruguay, between 1 March and 31 October 2006.

Patients had rectal swabs obtained 1, 4, 7, 10, 13 and 16 days after admission. Samples were streaked on two MacConkey agar plates supplemented with 4 mg/L ceftazidime or 2 mg/L ciprofloxacin, respectively. Putative enterobacterial colonies were identified as previously described [9]. Only the first antibiotic-resistant isolate per patient was included in the study.

2.2. Bacterial strains

Fifty-six fluoroquinolone- and/or oxyimino-cephalosporin-resistant enterobacterial isolates were studied. Strains *Klebsiella pneumoniae* 120, *E. coli* 137 and *K. pneumoniae* 2296 (provided by J. Martinez and J. Calvo Montes) were used as positive controls for *qnrA*, *qnrB* and *qnrS*, respectively.

2.3. Antibiotic susceptibility testing and extended-spectrum β -lactamase screening

Antibiotic susceptibility was determined by disk diffusion on Mueller–Hinton agar and was interpreted according to Clinical and Laboratory Standards Institute (CLSI) recommendations [10]. Tested antibiotics included ampicillin, amoxicillin/clavulanic acid, cefalothin, cefotaxime, ceftazidime, meropenem, cefoxitin, ciprofloxacin, cefuroxime, gentamicin, amikacin, cefepime and imipenem (Oxoid Ltd., Basingstoke, UK).

All isolates underwent ESBL detection by comparative testing using ceftazidime/clavulanic acid and cefotaxime/clavulanic acid disks (30/10 μ g) (Oxoid Ltd.) with plain ceftazidime and cefotaxime disks according to CLSI recommendations [10].

The ciprofloxacin minimum inhibitory concentration was determined by Etest following CLSI guidelines [10].

2.4. Extended-spectrum β -lactamase detection

Positive ESBL screening results were further analysed by polymerase chain reaction (PCR) for the presence of *bla*_{CTX-M} using

generic [11] and specific primers for the groups *bla*_{CTX-M-1} [8], *bla*_{CTX-M-2} [6] and *bla*_{CTX-M-9} (CTX-M-9F, 5'ATGGTGACAAAGAGAR-TGCAA; and CTX-M-9R, 5'TTACAGCCCTTCGGCGATGAT). *bla*_{TEM}, *bla*_{PER-2} and *bla*_{SHV} genes were also sought using specific primers [9,11]. PCR products were sequenced using the same primers.

2.5. *qnr* gene screening

The genes *qnrA*, *qnrB* and *qnrS* were sought by means of multiplex PCR and sequencing using primers previously described by Robicsek et al. [12].

2.6. Characterisation of class 1 integrons

All 56 isolates were analysed for the presence of class 1 integrons by PCR using primers I5/I3 [6]. Positive isolates were further studied looking for the variable region, the 3'-conserved region and ISCR1 using primers 5'CS/3'CS, *qacE1F*/*sul1b* and ORFend/F12R, respectively [6].

The region located between ISCR1 and the second *qacE Δ 1-sul1* was amplified with primers *orf513InK13/qacE1b* and the PCR product was sequenced using a gene walking strategy as reported previously [6]. The region between *qnrA* and the second 3'CS structure was elucidated with the primer *qnrAFend* (5'AGTAATAGTGCTTCCGGATTA), as shown in Fig. 1A.

2.7. Detection and mapping of class 2 integrons

Class 2 integrase was amplified by PCR and was sequenced employing primers *IntI2.F/IntI2.R* as described by Marquez et al. [7] in order to include the stop codon usually present in this gene. The genetic arrangement was determined by PCR mapping and sequencing of PCR products using specific primers for the genes *aadA1* (*aadA1F*, 5'GGCAGCCAATGACATTCTT; *aadA1R*, 5'CCAAGC-GATCTTCTTGCCAA), *dfr1* [13], *sat1* (*sat1F*, 5'CTATCTACCAGAA-GTGTGAGC; *sat1R*, 5'GTTTCGTTCCGAGACITGAGG) and *orfX* (*orfXA*, 5'GCGATCTGTTGAAGGTGGTT; *orfXB*, 5'GATACAT-GATCTTCAGGCC). Primers for *sat1* and *orfX* were kindly pro-




Fig. 1. Diagram depicting the different (A) class 1 and (B) class 2 integrons detected in this work. Roman numerals in (A) and lower case letters in (B) indicate the type of genetic arrangement. Small black arrows represent the different primers used to map the integrons; their location is relative to the DNA sequence to which they hybridise. Filled ovals represent the 59-bp element. Black boxes indicate the *intI1* promoter region and the *attI* site. Asterisks in (B) depict the internal stop codon usually present in these integrons. As shown in genetic arrangement c, the internal stop codon has disappeared, whereas the rest of this integron could not be characterised (symbolised by a striped box).

vided by Dr Pablo Power (Facultad de Farmacia y Bioquímica, Buenos Aires, Argentina). The strategy used for gene walking is shown in Fig. 1B.

2.8. Pulsed-field gel electrophoresis (PFGE)

Resistant microorganisms acquired after admission to the ICU harbouring any of the genes under study and suspected of belonging to an episode of clonal dissemination were analysed by PFGE as previously reported [9].

2.9. Replicon typing

Strains carrying ESBLs and/or *qnr* genes underwent PCR replicon typing according to Carattoli et al. [14] to determine the incompatibility groups of plasmids carrying such genes.

3. Results

In total, 56 isolates showed resistance to ceftazidime and/or ciprofloxacin (24 *Enterobacter* spp., 16 *E. coli*, 12 *K. pneumoniae* and 4 *Citrobacter freundii*), of which 29 carried at least one of the genes under study (Table 1). Eleven isolates (six *K. pneumoniae*, one *E. coli*, three *Enterobacter* spp. and one *C. freundii*) yielded positive results in presumptive and confirmatory ESBL detection tests.

Five isolates (three *C. freundii* and two *Enterobacter cloacae*) harboured *qnr* alleles. Twenty-four strains presented either class 1 or 2 integrons. Table 1 depicts the main features of the 29 strains.

Of the 56 colonised patients, 29 already carried resistant microorganisms upon admission into the ICU, whereas the remaining 27 acquired resistant microorganisms after admission.

Furthermore, 16 (28.6%) of the 56 patients had the same drug-resistant microorganisms (at the species level and susceptibility profile) isolated either from the gut and/or pharynx as well as from diverse clinical samples (e.g. peritoneal fluid, surgical wound, tracheal aspirate, blood culture, urine culture, etc.) (data not shown). Ten of the sixteen microorganisms isolated from the gut or pharynx of such patients carried at least one of the genes under study (Table 1).

Unfortunately, most of the information regarding clinical samples was obtained retrospectively; hence, microorganisms isolated from such samples were no longer available for molecular typing assays.

3.1. Extended-spectrum β -lactamase identification

CTX-M-derived enzymes were detected in 10 of 11 ESBL-producing isolates; six (*K. pneumoniae*) corresponded to CTX-M-2, one (*E. coli*) to CTX-M-15 and three (two *E. cloacae* and one *Enterobacter aerogenes*) to CTX-M-9. In addition, PER-2 was detected in one *C. freundii* isolate. No TEM- or SHV-derived ESBL were detected in these strains.

3.2. Screening and determination of *qnr* allelic variants

Two *E. cloacae* isolates (4R-HC112 and 2R-HC158) and three *C. freundii* isolates (1R-HC157, 1R-HC164 and 1R-HC175) harboured *qnr* variants. These isolates were obtained from different ICU inpatients; interestingly, all microorganisms carrying *qnrB* alleles were isolated upon admission to the unit. Both *E. cloacae* isolates harboured *qnrA1*, whereas the *C. freundii* isolates presented three different *qnrB* variants, namely *qnrB4* (1R-HC175), *qnrB13* (1R-HC157) and *qnrB17* (1R-HC164). All of these isolates remained susceptible to ciprofloxacin (Table 1).

Surprisingly, whereas *qnrA1*-carrying isolates displayed a multiresistant profile [including resistance to

trimethoprim/sulfamethoxazole (SXT), gentamicin and oxyimino-cephalosporins], those carrying *qnrB13* and *qnrB17* alleles were only resistant to oxyimino-cephalosporins.

3.3. Characterisation of class 1 integrons

Class 1 integrons were detected in 21 of the 56 studied strains (Table 1), clustered in six different genetic arrays. Four such arrays were simple integrons and the other two were complex integrons carrying the ISCR1 element.

Apart from *sul1*, responsible for resistance to sulfonamides, eight different antibiotic resistance gene cassettes were found within the six arrays, namely resistance to aminoglycosides [*aadA1*, *aadA2*, *aadA5*, *aadB* and *aac(6')-Ib*], β -lactams (*bla_{OXA-2}* and *bla_{OXA-9}*) and trimethoprim (*dfrA17*).

The complex class 1 integrons showed the following genes linked to ISCR1: *bla_{CTX-M-2}* (resistance to oxyimino-cephalosporins); and *qnrA1* (diminished susceptibility to fluoroquinolones) and the regulator of class C β -lactamases, *ampR* (genetic arrangements VI and V, respectively; see Fig. 1A).

bla_{CTX-M-2} was always found embedded in a complex class 1 integron, with a variable region formed by the cassettes *aac(6')-Ib-bla_{OXA-2}-orfD*. This particular complex class 1 integron confers resistance to aminoglycosides (amikacin) and oxyimino-cephalosporins, two major antibiotic groups used within ICUs. The arrangement *aac(6')-Ib-bla_{OXA-2}-orfD* was detected in six isolates, with this arrangement being second in frequency to variable regions composed of a single gene cassette, namely *aadA1*, present in nine strains (Table 1).

Another relevant integron is that identified as arrangement V (Fig. 1A), whose variable region is constituted by the genes *aadB-aadA2*, whereas the genes associated with ISCR1 are *qnrA1-ampR*. This integron confers resistance to gentamicin (mediated by *aadB*) as well as diminished susceptibility to fluoroquinolones on account of *qnrA1*; this integron was found to be encoded in a conjugative plasmid >50 kb that also carries the gene *bla_{CTX-M-9}* (data not shown).

3.4. Characterisation of class 2 integrons

Class 2 integrons were detected in 13 isolates. Interestingly, most of these isolates (10/13) also carried class 1 integrons.

The genetic arrangement *dfrA1-sat2-aadA1-orfX* was found in 12 class 2 integrons. Although this particular arrangement does not confer resistance to any antibiotic of clinical use, the simultaneous presence of class 1 and 2 integrons in a bacterial strain confers resistance to SXT. Ten such class 2 integrons had the IS5 element located between the genes *intI2* and *dfrA1* (Fig. 1B). To our understanding, this is the first report of an IS5 element inserted in a class 2 integron. This insertion sequence has been implicated in loss of porins and the emergence of resistance to carbapenems [15]. Considering that the gene cassette promoter in class 2 integrons is not clearly defined, the presence of this insertion sequence upstream of such cassettes could contribute not only to their horizontal transfer but also their expression.

Finally, one of the class 2 integrons (isolate 1R-HC081) lacked the stop codon (amino acid position 174) usually present in *intI2* genes. In this particular case, we were unable to determine the variable region, regardless of the various combinations of primers used. More studies are necessary to elucidate the genes encoded in this integron.

3.5. Pulsed-field gel electrophoresis patterns

Six *K. pneumoniae* and eight *E. cloacae* isolates were analysed by PFGE. Five of the *K. pneumoniae* isolates were grouped in

Table 1
Main features of isolates carrying extended-spectrum β -lactamases (ESBLs), *qnr* alleles, and class 1 and 2 integrons.

| Sample no. ^a | LoS (days) ^b | Microorganism | Class 1 integron | | Class 2 integron | | ESBL | <i>qnr</i> | PFGE pulsotype | Inc. group ^c | CIP MIC (μ g/mL) | Resistance profile |
|-------------------------|-------------------------|---------------------------------|------------------|---------------------|------------------|---------------------|----------|---------------|----------------|-------------------------|-----------------------|------------------------|
| | | | <i>int1</i> | Genetic arrangement | <i>int2</i> | Genetic arrangement | | | | | | |
| 2R-HC080 ^d | 25 | <i>Citrobacter freundii</i> | - | - | - | - | PER-2 | - | - | P, F | 1 | TGC, FQ, GEN, SXT |
| 1R-HC081 | 10 | <i>Escherichia coli</i> | - | - | + | c | - | - | - | - | 6 | FQ, SXT |
| 1R-HC093 | 6 | <i>E. coli</i> | + | III | - | - | - | - | - | - | 8 | TGC, FQ, GEN, SXT |
| 1R-HC094 ^d | 27 | <i>Enterobacter cloacae</i> | + | I | + | a | - | - | - | - | >32 | TGC, FQ, SXT |
| 2F-HC096 | 22 | <i>Klebsiella pneumoniae</i> | + | VI | - | - | CTX-M-2 | - | B1 | F, A/C | 0.064 | TGC, GEN, SXT |
| 2F-HC109 ^d | 47 | <i>E. cloacae</i> | + | I | - | - | - | - | 3c | - | >32 | TGC, FQ, AMK, GEN, SXT |
| 2R-HC110 | 13 | <i>K. pneumoniae</i> | + | II | + | a | - | - | C | - | >32 | TGC, FQ, AMK, SXT |
| 4R-HC112 | 13 | <i>E. cloacae</i> | + | V | - | - | CTX-M-9 | <i>qnrA1</i> | 1 | H11, H12 | 0.38 | TGC, GEN, SXT |
| 1F-HC113 ^d | 9 | <i>Enterobacter aerogenes</i> | - | - | - | - | CTX-M-9 | - | - | K | 0.023 | TGC, GEN, SXT |
| 1R-HC114 | 7 | <i>E. coli</i> | + | IV | + | a | - | - | - | - | >32 | TGC, FQ, AMK, GEN, SXT |
| 1R-HC124 | 9 | <i>E. cloacae</i> | + | II | + | b | - | - | - | - | >32 | TGC, FQ, AMK, GEN, SXT |
| 3R/F-HC125 | 8 | <i>K. pneumoniae</i> | + | VI | - | - | CTX-M-2 | - | B2 | F, A/C | 0.023 | TGC, GEN, SXT |
| 3R-HC136 ^d | 60 | <i>E. cloacae</i> | + | I | + | a | - | - | 3d | - | >32 | TGC, FQ, SXT |
| 3R/F-HC146 ^d | 16 | <i>E. cloacae</i> | + | I | + | a | - | - | 2b | - | >32 | TGC, FQ, AMK, GEN, SXT |
| 5R-HC149 ^d | 19 | <i>E. cloacae</i> | + | I | + | a | - | - | 2a | - | >32 | TGC, FQ, AMK, GEN, SXT |
| 1R/F-HC154 ^d | 2 | <i>E. cloacae</i> | - | - | + | b | - | - | - | - | >32 | TGC, FQ, SXT |
| 1R-HC152 | 42 | <i>K. pneumoniae</i> | + | VI | - | - | CTX-M-2 | - | - | A/C | >32 | TGC, FQ, AMK, GEN, SXT |
| 1R-HC151 ^d | 66 | <i>E. cloacae</i> | + | I | + | a | - | - | - | - | >32 | TGC, FQ, AMK, GEN, SXT |
| 1R-HC157 | 41 | <i>C. freundii</i> | + | - | - | - | - | <i>qnrB13</i> | - | B/O, K | 0.125 | TGC |
| 2R-HC158 | 11 | <i>E. cloacae</i> | + | V | - | - | CTX-M-9 | <i>qnrA1</i> | 1 | H11, H12 | 0.25 | TGC, GEN, SXT |
| 2R-HC159 | 14 | <i>E. coli</i> | - | - | - | - | CTX-M-15 | - | - | F | >32 | TGC, FQ, GEN, SXT |
| 6R-HC160 | 17 | <i>K. pneumoniae</i> | + | VI | - | - | CTX-M-2 | - | A | A/C | >32 | TGC, FQ, GEN, SXT |
| 2R-HC161 | 8 | <i>E. cloacae</i> | + | I | + | a | - | - | 3a | - | >32 | TGC, FQ, AMK, GEN, SXT |
| 1R-HC164 | 6 | <i>C. freundii</i> | - | - | - | - | - | <i>qnrB17</i> | - | K | 0.125 | TGC |
| 5R-HC167 | 17 | <i>K. pneumoniae</i> | + | VI | - | - | CTX-M-2 | - | A | A/C | >32 | TGC, FQ, AMK, GEN |
| 6R-HC168 ^d | 29 | <i>K. pneumoniae</i> | + | VI | - | - | CTX-M-2 | - | A | A/C | >32 | TGC, FQ, AMK, GEN |
| 4R-HC170 | 17 | <i>E. cloacae</i> | + | I | + | a | - | - | 3b | - | >32 | TGC, FQ, AMK, GEN, SXT |
| 1R-HC174 | 12 | <i>Enterobacter intermedium</i> | + | I | - | - | - | - | - | - | >32 | TGC, FQ, AMK, GEN, SXT |
| 1R-HC175 | 1 | <i>C. freundii</i> | - | - | + | a | - | <i>qnrB4</i> | - | K | 0.25 | TGC, GEN, SXT |

PFGE, pulsed-field gel electrophoresis; CIP, ciprofloxacin; MIC, minimum inhibitory concentration; TGC, third-generation cephalosporins; FQ, fluoroquinolones; GEN, gentamicin; SXT, trimethoprim/sulfamethoxazole; AMK, amikacin.

^a Numerals indicate the number of the sample that yielded the first antibiotic-resistant isolate; R/F indicates whether the sample was of rectal or pharyngeal origin.

^b Length of intensive Care Unit stay.

^c Plasmid incompatibility group determined by replicon typing.

^d Patients in which the same microorganism (species level and susceptibility profile) was obtained from clinical samples and rectal/pharyngeal swabbing.




Fig. 2. Pulsed-field gel electrophoresis (PFGE) analysis of nosocomial isolates of *Klebsiella pneumoniae* and *Enterobacter cloacae*. Lanes 1–4, pulsed types of *K. pneumoniae* (lane 1, pulsed type A; lane 2, pulsed type B1; lane 3, pulsed type B2; and lane 4, pulsed type C); lanes 5 and 13, *Salmonella* serotype Braenderup; lanes 6–12, pulsed types of *E. cloacae* (lane 6, pulsed type 1; lane 7, pulsed type 2a; lane 8, pulsed type 2b; lane 9, pulsed type 3a; lane 10, pulsed type 3b; lane 11, pulsed type 3c; and lane 12, pulsed type 3d).

two clusters: pulsed type A (6R-HC160, 5R-HC167 and 6R-HC168), and subtypes B1 (2F-HC096) and B2 (3R/F-HC125) (Table 1; Fig. 2). Whilst isolates belonging to pulsed type A were resistant to ciprofloxacin, those belonging to pulsed type B were susceptible to this antibiotic.

On the other hand, *E. cloacae* isolates 2R-HC158 and 4R-HC112 (carrying *bla*_{CTX-M-9} and *qnrA1*) shared the same pulsed type (i.e. pulsed type 1). Interestingly, they showed no resemblance to pulsed types 2 or 3, which did not harbour ESBLs.

3.6. Replicon typing

Isolates carrying ESBL genes were linked to various incompatibility groups. In this regard, those bearing *bla*_{CTX-M-2} were associated with groups A/C, those carrying *bla*_{CTX-M-9} belonged to groups HI and K, and the isolate carrying *bla*_{CTX-M-15} belonged to group F. On the other hand, plasmids bearing *qnrA* belonged to group HI, whereas those harbouring the allelic variants of *qnrB* belonged to group K (Table 1).

4. Discussion

This is the first report regarding the situation of CTX-M ESBLs within an ICU in Uruguay. Despite the small number of samples, the results show notable differences with those obtained in Argentina [4] and with a previous report from Uruguay in which the detected enzymes corresponded only to the CTX-M-2 group [5,6].

Although CTX-M-2 is the most frequently found enzyme in this report (6/11), enzymes CTX-M-9 and CTX-M-15 constitute an important fraction (4/11) of ESBL-producing strains.

The variants detected in Uruguay belong to the same group of ESBLs described recently by Pallecchi et al. in Peru and Bolivia [2]. The current results could be compatible with the observations made by these authors regarding the dissemination and expansion of this group of β -lactamases in Latin America.

The ESBL diversity detected in this work probably constitutes a fraction of the actual number of enzymes circulating within the ICU under study. Considering the relatively high concentration of ceftazidime (4 mg/L) used in this work to select resistant microorganisms, strains harbouring ESBLs with weak activity against this

drug (such as some variants of the CTX-M enzymes) may have been missed.

In this work, we also report for the first time the detection of *qnr* genes in Uruguay. *qnrA* alleles were found to be inserted in a complex class 1 integron with a genetic arrangement undescribed so far: *intI1-aadB-aadA2-qacE Δ 1-sul1::ISCR1::qnrA1::ampR*. To our understanding, this constitutes the first description of the genomic context of *qnrA1* in Latin America.

Regarding the occurrence of class 1 and 2 integrons, an important presence of genes conferring resistance to broad-spectrum antibiotics such as amikacin [*aac(6')-Ib*], gentamicin (*aadB*), oxyimino-cephalosporins (*bla*_{CTX-M-2}) and quinolones (*qnrA1*) was found. This differs from a recent report by Marquez et al. [5] regarding *E. coli* isolates causing urinary tract infections in outpatients. Their report describes most gene cassettes as variants of *aadA* and *dfrA*, which confer resistance to streptomycin and trimethoprim, respectively. The extensive presence of *aadA* in both settings could be due to gratuitous selection on account of its widespread presence in class 1 integrons.

Irrespective of this, the difference in the amount of antibiotic resistance genes in integrons in isolates from hospital and community settings could be explained by the selective pressure exerted by the different antibiotics present in such environments.

On the other hand, the integron carrying *bla*_{CTX-M-2} is of particular interest since it has been stably present in our region for over a decade [6].

Finally, the presence in an ICU of a class 2 integron with a presumptively functional integrase could set the stage for recombination events and acquisition of new gene cassettes by this genetic structure. So far there have been few reports describing such class 2 integrases, but only strains of human origin were linked to antibiotic resistance genes (*dfrA14*) [7].

The presence of microorganisms with a functional *intI2* gene within an ICU where most class 1 integrons either code for *aac(6')-Ib* or *aadB* could imply that it is only a matter of time before we find class 2 integrons carrying those gene cassettes, thus further contributing to their horizontal dissemination.

Funding: This work was partially supported by grants from S/C/OP/76/30 PDT (Programa de Desarrollo Tecnológico, Ministerio de Educación y Cultura del Uruguay) to RV as well as grants

from CSIC (Comisión Sectorial de Investigación Científica, Uruguay) to NFC. Part of this work was also supported by grants LSHM-CT-2003-503335 from the European Community and BFU2006-04574 from Ministerio de Ciencia e Innovación, Spain, to JAA and grants from UBACYT and ANPCYT to GG.

Competing interests: None declared.

Ethical approval: Not required.

References

- [1] Canton R, Coque TM. The CTX-M β -lactamase pandemic. *Curr Opin Microbiol* 2006;9:466–75.
- [2] Pallecchi L, Bartoloni A, Fiorelli C, Mantella A, Di Maggio T, Gamboa H, et al. Rapid dissemination and diversity of CTX-M extended-spectrum β -lactamase genes in commensal *Escherichia coli* isolates from healthy children from low-resource settings in Latin America. *Antimicrob Agents Chemother* 2007;51:2720–5.
- [3] Strahilevitz J, Jacoby GA, Hooper DC, Robicsek A. Plasmid-mediated quinolone resistance: a multifaceted threat. *Clin Microbiol Rev* 2009;22:664–89.
- [4] Quinteros M, Radice M, Gardella N, Rodriguez MM, Costa N, Korbenfeld D, et al. Extended-spectrum β -lactamases in Enterobacteriaceae in Buenos Aires, Argentina, public hospitals. *Antimicrob Agents Chemother* 2003;47:2864–7.
- [5] Marquez C, Labbate M, Raymondo C, Fernandez J, Gestal AM, Holley M, et al. Urinary tract infections in a South American population: dynamic spread of class 1 integrons and multidrug resistance by homologous and site-specific recombination. *J Clin Microbiol* 2008;46:3417–25.
- [6] Vignoli R, Cordeiro N, Seija V, Schelotto F, Radice M, Ayala J, et al. Genetic environment of CTX-M-2 in *Klebsiella pneumoniae* isolates from hospitalized patients in Uruguay. *Rev Argent Microbiol* 2006;38:84–8 [in Spanish].
- [7] Marquez C, Labbate M, Ingold AJ, Chowdhury PR, Ramirez MS, Centron D, et al. Recovery of a functional class 2 integron from an *Escherichia coli* strain mediating a urinary tract infection. *Antimicrob Agents Chemother* 2008;52:4153–4.
- [8] Cordeiro NF, Robino L, Medina J, Seija V, Bado I, Garcia V, et al. Ciprofloxacin-resistant enterobacteria harboring the *aac(6)-Ib-cr* variant isolated from feces of inpatients in an intensive care unit in Uruguay. *Antimicrob Agents Chemother* 2008;52:806–7.
- [9] Vignoli R, Calvelo E, Cordeiro NF, Lucero R, Ingold E, Quintana A, et al. Association of broad-spectrum antibiotic use with faecal carriage of oxyiminocephalosporin-resistant Enterobacteriaceae in an intensive care unit. *J Hosp Infect* 2006;63:306–15.
- [10] Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing. Nineteenth informational supplement. Document M100-S19. Wayne, PA: CLSI; 2009.
- [11] Yang D, Guo Y, Zhang Z. Combined porin loss and extended spectrum β -lactamase production is associated with an increasing imipenem minimal inhibitory concentration in clinical *Klebsiella pneumoniae* strains. *Curr Microbiol* 2009;58:366–70.
- [12] Robicsek A, Strahilevitz J, Sahn DF, Jacoby GA, Hooper DC. *qnr* prevalence in ceftazidime-resistant Enterobacteriaceae isolates from the United States. *Antimicrob Agents Chemother* 2006;50:2872–4.
- [13] Levings RS, Lightfoot D, Partridge SR, Hall RM, Djordjevic SP. The genomic island SG11, containing the multiple antibiotic resistance region of *Salmonella enterica* serovar Typhimurium DT104 or variants of it, is widely distributed in other *S. enterica* serovars. *J Bacteriol* 2005;187:4401–9.
- [14] Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 2005;63:219–28.
- [15] Song W, Suh B, Choi JY, Jeong SH, Jeon EH, Lee YK, et al. In vivo selection of carbapenem-resistant *Klebsiella pneumoniae* by OmpK36 loss during meropenem treatment. *Diagn Microbiol Infect Dis* 2009;65:447–9.